## PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through the skin

wear long-sleeved shirt, long pants, shoes, socks, and waterproof gloves. Any from children, domestic animals and pets. All handlers (including applicators) must must wear waterproof gloves. person who retrieves carcasses or unused bait following application of this product Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep away

drinking, chewing gum, using tobacco or using the toilet, and change into clean As soon as possible, wash hands thoroughly after applying bait and before eating hot water. Keep and wash PPE separately from other laundry. Remove PPE cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keen and useh DDE account. Communications for washables, use detergent and immediately after handling this product. Wash the outside of gloves before removing.

Have this label with you when obtaining treatment advice.

advice. Have person sip a glass of water if able to swallow. Do not induce vomiting If swallowed: Call a poison control center or doctor immediately for treatment unless told to do so by the poison control center or doctor.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Call a poison control center or doctor immediately for treatment advice.

treatment advice. water for 15-20 minutes. Call a poison control center or doctor immediately for If on skin or clothing: Take off contaminated clothing. Rinse skin with plenty of cool

If swallowed, this material may reduce the clotting ability of blood and cause bleeding normal (up to 30 days). In severe cases, blood transfusions may be needed. Also for pets, if needed, check prothrombin time every 3 days until values return to (bleeding or prolonged prothrombin times), give Vitamin  $K_1$  intramuscularly or orally, For humans or animals ingesting bait and/or showing obvious poisoning symptoms NOTE TO PHYSICIAN OR VETERINARIAN: Contains the anticoagulant difethialone. TREATMENT FOR PET POISONING: If animal eats bait, call veterinarian at once

mean high water mark. Do not contaminate water when disposing of equipment poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the birds. Dogs, cats and other predatory and scavenging mammals and birds might be ENVIRONMENTAL HAZARDS: This product is extremely toxic to mammals and

# STORAGE AND DISPOSAL

Pesticide Storage: Store in original container in a cool, dry place inaccessible to Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed children and pets.

of in trash or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. and dispose of in a sanitary landfill. Completely empty container, then offer for recycling or reconditioning; or puncture

Buyer assumes all risk of use and/or handling of this material when such use and/or expressed or implied, concerning the use of this product other than indicated on the label WARRANTY: To the extent consistent with applicable law, seller makes no warranty handling is contrary to label instructions.

EPA Est. No. 7173-WI-1 EPA Reg. No. 7173-218



## ALL WEATHER BAIT FOR WET OR DRY AREAS FOR INDOOR AND OUTDOOR USE

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN-RESISTANT HOUSE MICE

NORWAY RATS AND HOUSE MICE MAY CONSUME FOUR OR FIVE DAYS AFTER FEEDING BEGINS A LETHAL DOSE IN ONE NIGHT'S FEEDING WITH FIRST DEAD RODENTS APPEARING

Total	Inert Ingredients	Activ
	=	0
	Q	3
	7	9
	~	7
	=	8
•	Œ	-
	=	9
	S	=
-	-	
	-	$\Omega$
		7
		1
	-	3
•	-	a
	•	5
•	-	=
	•	0
	0	
	-	
-		
-	-	
•		•
	•	
•	ē .	
	•	
_	-	•
Total 100.0000%	. 99.9975%	Active Ingredient: difethialone 0.0025%

### AUTION See side panel for additional precautionary statements. KEEP OUT OF REACH OF CHILDREN



Liphatech, Inc. 3600 W. Elm Street Milwaukee, WI 53209 (800) 351-1476

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its

sites, pests, and application methods described on this label. READ THIS LABEL and follow all use directions and precautions. Use only for the

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides

- to help prevent accidents: Store product not in use in a location out of reach of children and pets.
- areas prone to vandalism. open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in secured or otherwise immobilized. Even stronger balt stations are needed in areas obtaining bait. If bait can be shaken from stations when they are lifted, units must be manner that prevents such children from reaching into bait compartments and to destruction by dogs and children under six years of age, and must be used in a nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant Apply bait in locations out of reach of children, pets, domestic animals, and
- on this label. Dispose of product container, unused, spoiled, and unconsumed bait as specified

the bait. bait stations must be used if children, pets, non-target mammals, or birds may access Note: Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant

crops. Do not broadcast bait. Burrow baiting with this product is prohibited. duct openings. Do not contaminate water, food, feedstuffs, food or feed handling containers holding less than 16 pounds of bait. Do not place near or inside ventilation structure, as defined above, is prohibited. Do not sell this product in individual associated with these sites. Fence and perimeter baiting beyond 100 feet from a aircraft), docks and port or terminal buildings, and related structures around and trash receptacles, agricultural and public buildings, transport vehicles (ships, trains or temporary residences, food processing facilities, industrial and commercial buildings rodent infestations. Examples of such structures include homes and other permanent so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting and house moe in and within 100 feet of man-made structures constructed in a manner USE RESTRICTIONS: This product can only be used to control Norway rats, roof rats equipment, or milk or meat handling equipment. Do not apply directly to food or feed

Remove as much alternative food as possible. or in locations where rats and/or house mice or their signs have been observed openings, in corners and concealed places, between floors or walls, beside burrows most likely find and consume bait. Generally, these are along walls, by gnawed Selection of Treatment Areas: Determine areas where rats and/or house mice will

APPLICATION DIRECTIONS: House Mice: Apply 1 block per placement, usually spaced 8 to 12 feet apart. Up to 2 blocks may be needed at points of very high house mouse activity, Maintain a constant supply of fresh bait for 15 days or until signs of house mouse activity cease.

a constant supply of fresh bait for 10 days or until signs of rat activity cease. Rats: Apply 6 to 23 blocks per placement, usually spaced 15 to 30 feet apart. Maintain

is not present, suspend bait a few inches above the high water line or place bait on a sewer grate, allowing just enough wire for bait to rest on manhole benching. If benching other end to a stationary structure such as the bottom step of manhole ladder or to a minimize chance for removal by rats or water. Place at least 23 blocks per manhole. board supported by opposing steps of the ladder. Securing bait in this manner will For sewer applications, securely attach one end of a wire to block of bait and the

stations and replenish bait as needed. does occur, repeat treatment. For a continuous infestation, set up permanent bai limit sources of rodent food, water and harborage as much as possible. If reinfestation bait that has become accessible to non-target animals. To discourage reinfestation gloves, collect and dispose of all visible dead animals and leftover bait, as well as any Follow-up: Replace contaminated or spoiled bait immediately. Wearing waterproof